NAME	MANUFACTURER	SOURCE	DELIVERY SYSTEM	TAKEN	ONSET, PEAK AND DURATION (APPROXIMATE HOURS)
apid-acting analogue					0 2 4 6 8 10 12 14 16 18 20 22 24 26 28 30 32 34 36
ovoRapid	Novo Nordisk	Analogue	Vial, cartridge, Novo prefilled pens, 1.6ml PumpCart	Just before / with / just after food	
umalog	Lilly	Analogue	Vial, cartridge, prefilled pen	Just before / with / just after food	
pidra	Sanofi	Analogue	Vial, cartridge, prefilled pen	0-15 mins before, or soon after, a meal	
hort-acting/neutral					
ctrapid	Novo Nordisk	Human	Vial	30 mins before food	
umulin S	Lilly	Human	Vial, cartridge	20–45 mins before food	
ypurin Bovine Neutral	Wockhardt UK	Bovine	Vial, cartridge	As advised by your healthcare team	
ypurin Porcine Neutral	Wockhardt UK	Porcine	Vial, cartridge	As advised by your healthcare team	
suman Rapid	Sanofi	Human	Cartridge	15–20 minutes before food	
fortion and law action					
Medium- and long-acting sulatard	Novo Nordisk	Human	Vial, cartridge, prefilled insulin doser	As advised by your healthcare team	
umulin l	Lilly	Human	Vial, cartridge, prefilled pen	As advised by your healthcare team As advised by your healthcare team	
ypurin Bovine Isophane	Wockhardt UK	Bovine	Vial, cartridge	As advised by your healthcare team As advised by your healthcare team	
ypurin Porcine Isophane	Wockhardt UK	Porcine	Vial, cartridge Vial, cartridge	As advised by your healthcare team As advised by your healthcare team	
purin Bovine Protamine Zinc	Wockhardt UK	Bovine	Vial Vial	As advised by your healthcare team As advised by your healthcare team	
ypurin Bovine Lente	THE RESERVE OF THE PERSON OF T	Bovine	Vial	As advised by your healthcare team As advised by your healthcare team	
suman Basal	Wockhardt UK Sanofi	Human	Vial, cartridge, prefilled pen	45–60 mins before food	
Sullidii Dasai	Salloll	Hulliali	viai, cartiluge, prefilieu peri	43-00 mins before food	
ixed					
umulin M3	Lilly	Human	Vial, cartridge, prefilled pen	20–45 mins before food	
ypurin Porcine 30/70 Mix	Wockhardt UK	Porcine	Vial, cartridge	As advised by your healthcare team	
suman Comb 15	Sanofi	Human	Cartridge	30–45 mins before food	
suman Comb 25	Sanofi	Human	Vial, cartridge, prefilled pen	30-45 mins before food	
suman Comb 50	Sanofi	Human	Cartridge	20–30 mins before food	
umalog Mix 25	Lilly	Analogue	Vial, cartridge, prefilled pen	Just before / with / just after food	
umalog Mix 50	Lilly	Analogue	Cartridge, prefilled pen	Just before / with / just after food	
lovoMix 30	Novo Nordisk	Anologue	Cartridge, prefilled pen	Just before / with / just after food	
ong-acting analogue					
antus	Sanofi	Analogue	Vial, cartridge, prefilled pen	Once a day, anytime (but at same time each day)	
vemir	Novo Nordisk	Analogue	Cartridge, prefilled pen, prefilled insulin doser	Once or twice a day, anytime (at same time each day)	
[resiba	Novo Nordisk	Analogue	Tresiba Penfill 100units/ml & prefilled Tresiba	Once a day, anytime (but preferably at the same time of the day)	
		400 000 and 500 500 500	FlexTouch 100units/ml & Tresiba FlexTouch		Tradition bounded 4th house
			200units/ml (NB the 3ml FlexTouch pen for		Tresiba, beyond 42 hours
			200units/ml contains 600 units)		

THE DIABETES UPDATE GUIDE TO MEDS & KIT, 2015

